

FIRST-ORDER LOGIC

CHAPTER 8

Pros and cons of propositional logic

- 😊 Propositional logic is **declarative**: pieces of syntax correspond to facts
- 😊 Propositional logic allows partial/disjunctive/negated information (unlike most data structures and databases)
- 😊 Propositional logic is **compositional**:
meaning of $B_{1,1} \wedge P_{1,2}$ is derived from meaning of $B_{1,1}$ and of $P_{1,2}$
- 😊 Meaning in propositional logic is **context-independent**
(unlike natural language, where meaning depends on context)
- 😞 Propositional logic has very limited expressive power
(unlike natural language)
E.g., cannot say “pits cause breezes in adjacent squares”
except by writing one sentence for each square

First-order logic

Whereas propositional logic assumes world contains **facts**,
first-order logic (like natural language) assumes the world contains

- **Objects**: people, houses, numbers, theories, Ronald McDonald, colors, baseball games, wars, centuries . . .
- **Relations**: red, round, bogus, prime, multistoried . . . ,
brother of, bigger than, inside, part of, has color, occurred after, owns,
comes between, . . .
- **Functions**: father of, best friend, third inning of, one more than, end of
. . .

Syntax of FOL: Basic elements

Constants *KingJohn, 2, UCB, ...*

Predicates *Brother, >, ...*

Functions *Sqrt, LeftLegOf, ...*

Variables *x, y, a, b, ...*

Connectives $\wedge \vee \neg \Rightarrow \Leftrightarrow$

Equality $=$

Quantifiers $\forall \exists$

Atomic sentences

Atomic sentence = *predicate*(*term*₁, ..., *term*_{*n*})
or *term*₁ = *term*₂

Term = *function*(*term*₁, ..., *term*_{*n*})
or *constant* or *variable*

E.g., *Brother*(*KingJohn*, *RichardTheLionheart*)
> (*Length*(*LeftLegOf*(*Richard*)), *Length*(*LeftLegOf*(*KingJohn*)))

Complex sentences

Complex sentences are made from atomic sentences using connectives

$$\neg S, \quad S_1 \wedge S_2, \quad S_1 \vee S_2, \quad S_1 \Rightarrow S_2, \quad S_1 \Leftrightarrow S_2$$

E.g. $Sibling(KingJohn, Richard) \Rightarrow Sibling(Richard, KingJohn)$

$$>(1, 2) \vee \leq(1, 2)$$

$$>(1, 2) \wedge \neg >(1, 2)$$

Truth in first-order logic

Sentences are true with respect to a **model** and an **interpretation**

Model contains ≥ 1 objects (**domain elements**) and relations among them

Interpretation specifies referents for


constant symbols \rightarrow **objects**

predicate symbols \rightarrow **relations**

function symbols \rightarrow **functional relations**

An atomic sentence $\textit{predicate}(\textit{term}_1, \dots, \textit{term}_n)$ is true
iff the **objects** referred to by $\textit{term}_1, \dots, \textit{term}_n$
are in the **relation** referred to by $\textit{predicate}$

Models for FOL: Example



Truth example

Consider the interpretation in which

Richard → Richard the Lionheart

John → the evil King John

Brother → the brotherhood relation

Under this interpretation, *Brother*(*Richard*, *John*) is true just in case Richard the Lionheart and the evil King John are in the brotherhood relation in the model

Models for FOL: Lots!

Entailment in propositional logic can be computed by enumerating models

We **can** enumerate the FOL models for a given KB vocabulary:

For each number of domain elements n from 1 to ∞

For each k -ary predicate P_k in the vocabulary

For each possible k -ary relation on n objects

For each constant symbol C in the vocabulary

For each choice of referent for C from n objects ...

Computing entailment by enumerating FOL models is not easy!

Universal quantification

$\forall \langle \text{variables} \rangle \langle \text{sentence} \rangle$

Everyone at Berkeley is smart:

$\forall x \text{ At}(x, \text{Berkeley}) \Rightarrow \text{Smart}(x)$

$\forall x \ P$ is true in a model m iff P is true with x being **each** possible object in the model

Roughly speaking, equivalent to the conjunction of instantiations of P

$$\begin{aligned} & (\text{At}(\text{KingJohn}, \text{Berkeley}) \Rightarrow \text{Smart}(\text{KingJohn})) \\ \wedge & (\text{At}(\text{Richard}, \text{Berkeley}) \Rightarrow \text{Smart}(\text{Richard})) \\ \wedge & (\text{At}(\text{Berkeley}, \text{Berkeley}) \Rightarrow \text{Smart}(\text{Berkeley})) \\ \wedge & \dots \end{aligned}$$

A common mistake to avoid

Typically, \Rightarrow is the main connective with \forall

Common mistake: using \wedge as the main connective with \forall :

$$\forall x \text{ } At(x, Berkeley) \wedge Smart(x)$$

means “Everyone is at Berkeley and everyone is smart”

Existential quantification

$\exists \langle variables \rangle \langle sentence \rangle$

Someone at Stanford is smart:

$\exists x \text{ } At(x, Stanford) \wedge Smart(x)$

$\exists x \text{ } P$ is true in a model m iff P is true with x being **some** possible object in the model

Roughly speaking, equivalent to the **disjunction** of instantiations of P

$$\begin{aligned} & (At(KingJohn, Stanford) \wedge Smart(KingJohn)) \\ \vee & (At(Richard, Stanford) \wedge Smart(Richard)) \\ \vee & (At(Stanford, Stanford) \wedge Smart(Stanford)) \\ \vee & \dots \end{aligned}$$

Another common mistake to avoid

Typically, \wedge is the main connective with \exists

Common mistake: using \Rightarrow as the main connective with \exists :

$$\exists x \text{ } At(x, Stanford) \Rightarrow Smart(x)$$

is true if there is anyone who is not at Stanford!

Properties of quantifiers

$\forall x \forall y$ is the same as $\forall y \forall x$ (why??)

$\exists x \exists y$ is the same as $\exists y \exists x$ (why??)

$\exists x \forall y$ is **not** the same as $\forall y \exists x$

$\exists x \forall y \text{ Loves}(x, y)$

“There is a person who loves everyone in the world”

$\forall y \exists x \text{ Loves}(x, y)$

“Everyone in the world is loved by at least one person”

Quantifier duality: each can be expressed using the other

$\forall x \text{ Likes}(x, \text{IceCream}) \quad \neg \exists x \neg \text{Likes}(x, \text{IceCream})$

$\exists x \text{ Likes}(x, \text{Broccoli}) \quad \neg \forall x \neg \text{Likes}(x, \text{Broccoli})$

Fun with sentences

Brothers are siblings

Fun with sentences

Brothers are siblings

$\forall x, y \text{ } Brother(x, y) \Rightarrow Sibling(x, y).$

“Sibling” is symmetric

Fun with sentences

Brothers are siblings

$$\forall x, y \text{ } Brother(x, y) \Rightarrow Sibling(x, y).$$

“Sibling” is symmetric

$$\forall x, y \text{ } Sibling(x, y) \Leftrightarrow Sibling(y, x).$$

One's mother is one's female parent

Fun with sentences

Brothers are siblings

$$\forall x, y \text{ Brother}(x, y) \Rightarrow \text{Sibling}(x, y).$$

“Sibling” is symmetric

$$\forall x, y \text{ Sibling}(x, y) \Leftrightarrow \text{Sibling}(y, x).$$

One's mother is one's female parent

$$\forall x, y \text{ Mother}(x, y) \Leftrightarrow (\text{Female}(x) \wedge \text{Parent}(x, y)).$$

A first cousin is a child of a parent's sibling

Fun with sentences

Brothers are siblings

$$\forall x, y \text{ Brother}(x, y) \Rightarrow \text{Sibling}(x, y).$$

“Sibling” is symmetric

$$\forall x, y \text{ Sibling}(x, y) \Leftrightarrow \text{Sibling}(y, x).$$

One's mother is one's female parent

$$\forall x, y \text{ Mother}(x, y) \Leftrightarrow (\text{Female}(x) \wedge \text{Parent}(x, y)).$$

A first cousin is a child of a parent's sibling

$$\forall x, y \text{ FirstCousin}(x, y) \Leftrightarrow \exists p, ps \text{ Parent}(p, x) \wedge \text{Sibling}(ps, p) \wedge \text{Parent}(ps, y)$$

Equality

$term_1 = term_2$ is true under a given interpretation
if and only if $term_1$ and $term_2$ refer to the same object

E.g., $1 = 2$ and $\forall x \neg (Sqrt(x), Sqrt(x)) = x$ are satisfiable
 $2 = 2$ is valid

E.g., definition of (full) *Sibling* in terms of *Parent*:

$$\forall x, y \text{ Sibling}(x, y) \Leftrightarrow [\neg(x = y) \wedge \exists m, f \neg(m = f) \wedge \\ \text{Parent}(m, x) \wedge \text{Parent}(f, x) \wedge \text{Parent}(m, y) \wedge \text{Parent}(f, y)]$$

Interacting with FOL KBs

Suppose a wumpus-world agent is using an FOL KB
and perceives a smell and a breeze (but no glitter) at $t = 5$:

$Tell(KB, Percept([Smell, Breeze, None], 5))$

$Ask(KB, \exists a \text{ Action}(a, 5))$

I.e., does KB entail any particular actions at $t = 5$?

Answer: $Yes, \{a/Shoot\}$ \leftarrow substitution (binding list)

Given a sentence S and a substitution σ ,

$S\sigma$ denotes the result of plugging σ into S ; e.g.,

$S = Smarter(x, y)$

$\sigma = \{x/Hillary, y/Bill\}$

$S\sigma = Smarter(Hillary, Bill)$

$Ask(KB, S)$ returns some/all σ such that $KB \models S\sigma$

Knowledge base for the wumpus world

“Perception”

$\forall b, g, t \text{ Percept}([Smell, b, g], t) \Rightarrow Smelt(t)$

$\forall s, b, t \text{ Percept}([s, b, Glitter], t) \Rightarrow AtGold(t)$

Reflex: $\forall t \text{ AtGold}(t) \Rightarrow \text{Action}(Grab, t)$

Reflex with internal state: do we have the gold already?

$\forall t \text{ AtGold}(t) \wedge \neg Holding(Gold, t) \Rightarrow \text{Action}(Grab, t)$

$Holding(Gold, t)$ cannot be observed

\Rightarrow keeping track of change is essential

Deducing hidden properties

Properties of locations:

$$\forall x, t \text{ } At(Agent, x, t) \wedge Smelt(t) \Rightarrow Smelly(x)$$

$$\forall x, t \text{ } At(Agent, x, t) \wedge Breeze(t) \Rightarrow Breezy(x)$$

Squares are breezy near a pit:

Diagnostic rule—infer cause from effect

$$\forall y \text{ } Breezy(y) \Rightarrow \exists x \text{ } Pit(x) \wedge Adjacent(x, y)$$

Causal rule—infer effect from cause

$$\forall x, y \text{ } Pit(x) \wedge Adjacent(x, y) \Rightarrow Breezy(y)$$

Neither of these is complete—e.g., the causal rule doesn't say whether squares far away from pits can be breezy

Definition for the *Breezy* predicate:

$$\forall y \text{ } Breezy(y) \Leftrightarrow [\exists x \text{ } Pit(x) \wedge Adjacent(x, y)]$$